

Application No. 10/826,743
Amendment dated May 1, 2006
Reply to Office Action of February 1, 2006

34

Docket No.: 60583(50530)

REMARKS

Claims 1-14 are pending. Applicants have amended claim 1. Support for the amendments can be found throughout the specification and in the claims as originally filed. Applicants make these amendments without prejudice to pursuing the original subject matter of this application in a later filed application claiming benefit of the instant application, including without prejudice to any determination of equivalents of the claimed subject matter. No new matter is introduced by these amendments.

Claims Rejections – 35 U.S.C. §112, second paragraph

Claims 1-5 and 9-14 are rejected, allegedly due to indefiniteness. Applicants traverse, but have amended claim 1 as follows. As an initial matter, the structures of Formulae (I) and (II) are replaced to comport with those delineated at page 4 of the specification as filed. Thus, what previously appeared as R_7 , R_8 in Formula (I) and R_3 and R_4 in Formula (II), now appears as R_5 and R_6 , and R_5 and R_6 in Formula (I) and (II), respectively. This is supported throughout the specification as filed including at page 4 and the examples.

Variable R_3 is removed in Formula (II) and replaced with R_5 . Support for this amendment appears throughout the specification as filed including at page 4.

The term "Each" which was previously capitalized in claim 1 is now amended to read "each".

Variables R_7 and R_8 are now delineated in claim 1. Support for this amendment appears throughout the specification as filed including at pages 3-6, and claims 1-3 as filed. As such, claims 2 and 3 no longer lack antecedent basis in claim 1, from which they both depend.

Application No. 10/826,743
Amendment dated May 1, 2006
Reply to Office Action of February 1, 2006

35

Docket No.: 60583(50530)

In regard to the rejection of claims 12-14 on the alleged basis that the nature of the recited agents are not known, Applicants traverse. Claims 12-14 recite anti-HCV agents. Applicants submit that one of ordinary skill would understand and appreciate the metes and bounds of an anti-HCV agent according to any of claims 12-14. For example, an anti-HCV agent is any agent that demonstrates HCV inhibitory activity, including those with demonstrated activity in standard assays such as those delineated in Examples 80 and 81 in the specification as filed. Such compounds include α -interferon, β -Interferon, ribavarin, and adamantine as delineated in the specification as filed, and also include agents known in the art, including those delineated in "HCV Anti-viral Agents", Griffith et al., *Annual Reports in Medicinal Chemistry*, Vol. 39, pp.223-237 (2004) (copy attached) and references cited therein, which describe various types of inhibitors, including those delineated in instant claims 12-14. As such, Applicants submit that claims 12-14 are not indefinite to one of ordinary skill in the relevant art and respectfully request withdrawal of the rejection.

Based on the foregoing amendments and reasoning, Applicants submit that the rejections should be withdrawn.

In view of the above, each of the presently pending claims in this application is believed to be in immediate condition for allowance. Accordingly, the Examiner is respectfully requested to pass this application to issue. Should any of the claims not be found to be in condition for allowance, the Examiner is requested to call Applicants' undersigned representative to discuss the application. Applicants thank the Examiner in advance for this courtesy.

Application No. 10/826,743
Amendment dated May 1, 2006
Reply to Office Action of February 1, 2006

36

Docket No.: 60583(50530)

The Director is hereby authorized to charge any credits or deficiency in the fees filed (or with any paper hereafter filed in this application by this firm) to our Deposit Account No. 04-1105, under Order No. (50530) 60583.

Dated: May 1, 2006

Respectfully submitted,

By

Jeffrey D. Hsi

Registration No.: 40,024

EDWARDS ANGELL PALMER & DODGE
LLP

P.O. Box 55874

Boston, Massachusetts 02205

(617) 439-4444

Attorneys/Agents For Applicant

HCV Anti-viral Agents

Ronald C. Griffith, Lillian Lou, Christopher D. Roberts, and Uli Schmitz

Genelabs Technologies Inc., 505 Penobscot Drive, Redwood City, CA 94063, USA

Contents	223
1. Introduction	224
2. Inhibitors of viral protein targets	224
2.1. NS3 protease	227
2.2. NS5b RNA dependent RNA polymerase	228
2.3. Non-nucleoside NS5b inhibitors	230
2.4. Nucleoside NS5b inhibitors	232
2.5. Other viral targets	232
3. Inhibitors for non-viral targets	232
3.1. Caspase	232
3.2. Toll-like receptor 7	233
3.3. Proteasome inhibitors	233
3.4. CD81	234
4. Conclusion	234
References	

1. INTRODUCTION

Chronic infection with HCV has become a major health problem associated with liver cirrhosis, hepatocellular carcinoma and liver failure. An estimated 170 million chronic carriers worldwide are at risk of developing liver disease [1]. In the US alone 2.7 million are chronically infected with HCV and the death-toll in 2000 was estimated between 8000 and 10,000, a number that is expected to increase significantly over the next years [2]. Considering the slow course of the infection and the nature of the current, insufficient therapy, treatment of HCV-related illness is a heavy burden on any public health system. Current therapy involving pegylated interferon- α alone (Pegasys, PEG-Intron™) or in combination with ribavirin (1) (Rebetrol™, Copegus™) [3], is expensive, associated with side-effects and exhibits only modest success rates, which are strongly dependent on the particular genotype of the virus. For the most prevalent HCV subtypes in the US, 1a and 1b (72%), the above co-therapy yields response rates between 40 and 60%. Note that the above therapy is not aimed at inhibiting viral targets directly, in contrast to the successful approaches in HIV therapy. Future therapies need to include the more elusive HCV subtypes and should benefit from targeting viral replication directly. HCV is a small, enveloped virus with a single strand RNA genome coding for a 3000 a.a. polyprotein which is processed into at least 10 individual viral proteins of which the six C-terminal ones are non-structural proteins termed NS2, NS3, NS4a, NS4b, NS5a and NS5b [4]. Assembly of the final

ANNUAL REPORTS IN MEDICINAL CHEMISTRY, VOLUME 39
ISSN: 0065-7743 DOI 10.1016/S0065-7743(04)39018-4

© 2004 Elsevier Inc.
All rights reserved


BEST AVAILABLE COPY

replication complex around the NS5b RNA polymerase requires five proteolytic steps of which the first is carried out by the metalloprotease NS2. Subsequently, the protease module of NS3 delivers the remaining cuts which most require NS4a as a cofactor for NS3. Fueled by the success of targeting the HIV protease and reverse transcriptase, the largest anti-HCV drug discovery efforts have been directed towards NS3 protease and NS5b RNA polymerase.

2. INHIBITORS OF VIRAL PROTEIN TARGETS

2.1. NS3 protease

NS3 is a relatively small serine protease of the chymotrypsin family and peptide sequence requirements for both efficient cleavage and competitive inhibition are well known [5]. Since low micromolar inhibition could be achieved with hexapeptides derived from N-terminal P-site cleavage products [6], the vast majority of work has not strayed from competitive peptidic inhibitors with either a covalent ('serine traps') or non-covalent mode of interaction. Most promising is the group of constrained tripeptides, 2–5, which arose from linking the P1 and P3 a.a. side chains and optimization of the P2 proline substituents [7]. Trailblazer BILN-2061, 2, yielded clinical proof-of-concept data as its outstanding *in vitro* activity in enzyme inhibition assays ($K_i = 0.3$ and 0.7 nM for subtype 1a and 1b, respectively) and the surrogate cell-based replicon assay ($EC_{50} = 3$ and 4 nM for subtype 1a and 1b, respectively) was mirrored by a dramatic 2–3 \log_{10} reduction of viral loads within 24 h in the majority of patients that had received 2 (200 mg in an oral solution of a PEG 400/ethanol 80:20 mixture, twice-daily for 2 days) [8]. While 2 was well tolerated up to 2000 mg and no serious clinical or laboratory findings were obtained in a human safety study, further clinical trials are currently on hold because cardiac lesions were observed in routine chronic safety testing in monkeys at supra-therapeutic doses [9,10]. The primary interaction site of BILN 2061 with NS3, as revealed by a crystal structure, involves largely the residues conserved across HCV subtypes [7]. This explains why K_i values for subtypes 2a, 2b and 3a are still below 100 nM. Oral bioavailability for 2 was below 20% in rats and rhesus monkeys; for dogs, values were as high as 38% [8]. Some of the clinical activity of 2 might in fact be the restoration of the IFN related cellular anti-viral response. NS3 has been shown to mediate



BEST AVAILABLE COPY

HCV A

inhibi
key a

Th
value:
by a c
of ma
and 1
aliph:
subst
vario
cyclo
repla
meml
used:
large
heter
[15].
carbo
exhib
non-f

Th
cente
boros
and c
mech
show
With
[18].
regio
NS3
[19].
seen
the c
with
meth
IC₅₀
17 [2
moie
cova
The
lacta
carb:
cryst
A15'
althc
chen


inhibition of the phosphorylation, hence activation of IFN regulatory factor-3 (IRF-3), a key anti-viral signaling molecule [11].

The scaffold of 2 tolerates small changes in several places. IC_{50} and replicon EC_{50} values of 3 and 4 are both below 10 nM [12]. When the carboxylate group in 3 is replaced by a cyclopropyl-acylsulfonamide moiety no loss in activity is observed. This latter class of macrocycles is further exemplified by 5, which exhibits IC_{50} and EC_{50} values of 21 nM and 124 nM, respectively [13]. Note that an unprotected amino group and a shortened aliphatic linker are compatible with good potency. Also unconstrained proline-substituted tripeptides, 6–10, retain activity. 6 represents a group of compounds wherein various bulky aliphatic substituents (e.g., *t*-butyl, cyclobutyl, cyclopentyl, and cyclohexyl) can be interchanged [13,14]. In this scaffold, the carboxylate group can be replaced by methyl- and phenyl-acylsulfonamide moieties. A wide range of six-membered and especially five-membered heterocycles, exemplified by 7 and 8, can be used in conjunction with the acylsulfonamide grouping [13]. 9 and 10 are examples of a large group of compounds with both IC_{50} and EC_{50} below 100 nM in which the heterocyclic proline-substituent assumes a range of substituted condensed ring systems [15]. Studies with acylsulfonamide bearing hexapeptides demonstrated that a cluster of carboxylate groups at the N-terminus contributes greatly to inhibitory activity; 11 exhibited an IC_{50} value of 3.8 nM [16]. Among the non-covalent NS3 inhibitors, two non-peptidic natural products, 12 and 13, showed submicromolar inhibition [17].


The class of covalent inhibitors is largely comprised of peptides bearing a reactive center at the cleavage site trapping the hydroxyl of Ser139, i.e., α -keto, aldehyde, boronic acid or lactam moieties. Most advanced is VX-950, 14, which binds slowly and covalently ($K_1^* = 3$ nM, $T_{1/2} = 58$ min), yet reversibly to NS3 with a multistep mechanism. The replicon EC_{50} is 400 nM and a cell-based viral clearance assay showed 3 log₁₀ reductions of viral RNA levels at concentrations above 3.5 μ M [18]. With good oral bioavailability and high liver exposure, 14 is slated for clinical studies [18]. The co-crystal structure of 14 shows that it only partially overlaps with the region accessed by BILN-2061, which explains why 14 retains full activity against NS3 mutants (e.g., D168V) that are more than 1000-fold less sensitive to BILN 2061 [19]. Modification of the bicyclopipeline moiety and the N-terminal capping group as seen in 15, yields potent derivatives ($IC_{50} < 500$ nM) [20]. Even the replacement of the cyclopentane moiety in 14 with the bulkier bicyclo[2.2.1]hept-2-ene is compatible with good potency [21]. Replacement of norvaline in 14 with difluoro-Abu and methylbenzyl instead of cyclopropyl yields potent anti-viral activity (replicon $IC_{50} = 630$ nM) [22]. Other examples of new α -keto-peptide inhibitors are 16 and 17 [23]. Compounds bearing only an aldehyde group replacing the entire α -keto-amide moiety also show *in vitro* activity in the low micromolar range [24]. Another class of covalent inhibitors is that of the pyrrolidine-*trans*-lactams, exemplified by 18 and 19. The crystal structure of a smaller example showed that the serine hydroxyl opens the lactam ring, forming a hemiketal [25]. As older compounds in this series exhibited carbamate moieties at the N-terminus, newer designs favor the urea linker. There is crystallographic evidence that both amide hydrogens engage in hydrogen bonding to A157 [26]. 18 and 19 exhibit replicon IC_{50} values of 300 and 100 nM, respectively, although 18 was found to be fivefold more active in the enzyme assay [26]. Due to the chemical instability of the lactam moiety, compounds like 18 and 19 exhibit

BEST AVAILABLE COPY


undesirable, fast intravenous (IV) clearance rates. GW0014, 20, however, although less potent, showed significantly improved *in vivo* pharmacokinetic properties and demonstrated anti-viral activity in a surrogate animal model with GBV-B infected marmosets following subcutaneous administration [27].



BEST AVAILABLE COPY



18 R



19 F

2.2. N

As an F
[28]. N
templat
terminu
compos
hydropl
solubili
 K_m for
truncat
purifica
case of
depend
Furt
withon

Table


K_m (P-
 K_m (U
IC₅₀ (t
His₆-t
Prime
IC₅₀ c

Smith et al.

although
ties and
infected

HCV Anti-viral Agents

227



2.2. NS5b RNA dependent RNA polymerase

As an RNA virus, HCV must encode its own RNA dependent RNA polymerase, NS5b [28]. NS5b is essential for viral replication and uses single-stranded HCV RNA as a template to initiate *de novo* synthesis [29]. The enzyme contains a hydrophobic C-terminus which anchors the polymerase to the ER membrane in a replicase complex composed of viral and host proteins [30]. Under enzymatic screening conditions the hydrophobic C-terminus of NS5b causes several important assay complications. The solubility of expressed protein is decreased by the presence of this tail as well as shifts in K_m for both RNA template and nucleotide substrates (Table 1). The usual solution is to truncate this tail as either the $\Delta 21$ or $\Delta 57$ construct. N- or C-terminal His₆ tags as purification aids can further change the affinities. This has the effect, particularly in the case of some non-nucleoside inhibitors, of causing IC₅₀ values to vary dramatically depending on the exact construct used (Table 1) [31].

Furthermore, the choice of homopolymeric vs. heteropolymeric templates with or without primers can further shift the substrate K_m s and consequently the inhibitor's

Table 1. Effect of NS5b enzyme constructs on kinetics and inhibition

	HT ^a -NS5b	HT-NS5b $\Delta 21$	NS5b $\Delta 57$ -HT	NS5b $\Delta 21$ -HT	NS5b
K_m (P-T) ^b (nM)	210	58	34	25	25
K_m (UTP) (nM)	6200	12,000	1800	5200	3300
IC ₅₀ (nM) ^c	54	440	2200	3000	5700


^a His₆-tagged.^b Primer-template.^c IC₅₀ of compound 21.

BEST AVAILABLE COPY

IC₅₀. Therefore one must clearly understand the enzymatic systems of reported results to properly analyze and interpret SAR. Crystal structures of the enzyme reveal similarities to the classic polymerase right-handed model with fingers, thumb and the active site palm subdomains [32,33]. Thus the unique structure, *de novo* mechanism and the absence of a homologous mammalian enzyme should allow potent and selective anti-virals to be discovered. Anti-viral therapy strategies targeting this polymerase have been approached with parallel efforts in both nucleoside and non-nucleoside inhibitor discovery.

2.3. Non-nucleoside NS5b inhibitors


Benzimidazole containing compounds were some of the first non-nucleoside inhibitors discovered to be active against NS5b and to date represent the most potent. **22** was reported to be active at <0.01 μ M [34] but under different enzyme assay conditions was reported at 0.28 μ M [35] demonstrating the importance of understanding the enzyme and polymer constructs. Similarly, the initially reported IC₅₀ of **21** using one enzyme construct was 0.054 μ M but using a different construct resulted in a 0.25 μ M IC₅₀ [31,35]. Kinetic studies show that they are non-competitive with NTP and competitive with the RNA template [31]. Further kinetic analysis demonstrated that **21** and **22** were exclusive inhibitors [36]. Finally, **22** was active in the HCV replicon (EC₅₀ = 0.35 μ M) and multiple resistant replicon mutants generated to **22** displayed a common P495L/A mutation in NS5b [35]. This proline residue is located in a recently reported allosteric GTP binding site [32] and may represent the binding site for this benzimidazole class of compounds. Further optimization of this class of allosteric inhibitors has yielded the optimized analog **23** (IC₅₀ < 0.5 μ M) [37–39] and the tetrazole **24** (IC₅₀ = 0.01 μ M) [40]. Interestingly, replacement of the benzimidazole with a pyrazolo-pyrimidine ring and reversal of its relative orientation (6:5 to 5:6 ring system) resulted in active compounds such as **25** (IC₅₀ < 1 μ M) [41].



R.C. Griffith *et al.*

systems of reported results of the enzyme reveal with fingers, thumb and the cture, *de novo* mechanism should allow potent and strategies targeting this both nucleoside and non-


(non-nucleoside inhibitors) the most potent. 22 was enzyme assay conditions nce of understanding the orted IC_{50} of 21 using one uct resulted in a $0.25 \mu M$ mpetitive with NTP and lysis demonstrated that 21 ive in the HCV replicon enerated to 22 displayed a ue is located in a recently the binding site for this of this class of allosteric $5 \mu M$) [37-39] and the ent of the benzimidazole rientation (6:5 to 5:6 ring μM) [41].




HCV Anti-viral Agents

BEST AVAILABLE COPY


229




A second allosteric binding site has been reported with inhibitors bound to crystalized NS5b [42,43]. Further optimization of one of these two compounds led to the discovery of 26 (NS5b $IC_{50} = 0.7 \mu M$, replicon $EC_{50} = 0.2 \mu M$). The crystal structure reveals the cyclohexyl ring binding tightly in a deep hydrophobic pocket [44]. Similarly, 27 binds to the same site with the cyclopentyl group binding to the same pocket. Recently, analogs of 27 were reported with activities less than $1 \mu M$ [45,46].




230

R.C. Griffith *et al.*


34



35



36



37

Other non-nucleoside inhibitors have been reported and characterized. A previously described benzothiadiazine (NS5b $IC_{50} = 0.08 \mu M$, replicon $EC_{50} = 0.5 \mu M$) was used to generate resistant replicons which mapped the key mutation to M414T, implicating a new inhibitor site on NS5b [47,48]. Similarly, a dimethoxypyrimidine analog 28 previously described [49] was used to generate resistant replicons containing either P156L or G152E mutations [47]. Interestingly, these mutations map the binding site of this compound to the active site, the first reported active-site non-nucleoside inhibitor. A non-nucleoside compound, HCV-371, was reported to have failed a phase I/II clinical trial [28,50]. While the compound was safe, it failed to reduce HCV RNA levels. However, follow-up compounds such as racemic 29 and (R)-pyranoidole 30 (NS5b $IC_{50} = 0.004$ and $0.06 \mu M$, respectively) have recently been reported [51,52]. An analog of dichloroacetamide 31, R803, is active in the replicon ($EC_{50} < 10 \mu M$) and is claimed to be an NS5b inhibitor [53]. Phase I results for this compound were reported and a phase I/II study scheduled to assess anti-viral efficacy [54]. Dihydropyrroles such as 32 (NS5b $IC_{50} < 5 \mu M$ for best compounds) and related pyrrolidine analogs such as 33 (NS5b $IC_{50} < 0.3 \mu M$ for best compounds) have been reported [55–57]. Triazine 34 (%inhibition at $0.1 \mu g/mL$: HCV NS5b = 46% and HBV RT = 52%) and difluorobenzamide 35 (%inhibition at $0.1 \mu g/mL$: HCV NS5b = 46% and HBV RT = 48%) are two reported promiscuous inhibitors with activity against both HCV NS5b and HBV RT [58,59]. Finally, compounds such as 36 (NS5b $IC_{50} < 10 \mu M$ for best compounds) [60] and 1,2-diaminophenyl compound 37 (NS5b $IC_{50} = 1.42 \mu M$) have recently been disclosed [61].

2.4. Nucleoside NS5b inhibitors

Nucleoside anti-virals are pro-drugs in that they are actively transported into cells and then activated by cellular kinases to the nucleotide triphosphate. This NTP is now able to competitively inhibit the enzyme or, more commonly, act as a substrate and be incorporated into the nascent RNA chain. Chemical or structural features of

HCV Ant

the inc
replicat
both) ti
inhibito
dioxola

To d
nucleos
 $IC_{50} =$
HCV r
nucleos
the NS
able to
substrat
with eit
were ge
this mu
group to
nucleoti
active i)
is curre
study (d
HCV R
kinetics
apparen




42

Recei
includin
base nu
45 were
for 45 d
with no

R.C. Griffith et al.

HCV Anti-viral Agents

231




37

ized. A previously
 $IC_{50} = 0.5 \mu M$) was
 tation to M414T,
 methoxypyrimidine
 resistant replicons
 , these mutations
 reported active-site
 was reported to
 as safe, it failed to
 cemic 29 and (R)-
 ave recently been
 ve in the replicon
 : I results for this
 anti-viral efficacy
 runds) and related
 ounds) have been
 NS5b = 46% and
 0.1 $\mu g/mL$: HCV
 is inhibitors with
 ounds such as 36
 nyl compound 37

ted into cells and
 NTP is now able
 substrate and be
 ural features of

the incorporated nucleoside subsequently prevents (chain-terminates) the further replication of the viral genome. While nucleoside inhibitors can vary in either (or both) the ribose or base portion of the molecule, the initially reported anti-HCV inhibitors were modified ribose analogs such as 2'-fluoro [62], 4'-azido [63] and dioxolane cytidine analogs [64].

To date, the most significant amount of work has been reported on 2'-methylribose nucleosides. Adenosine (38, $IC_{50} = 1.9 \mu M$, $EC_{50} = 0.17 \mu M$) and guanosine (39, $IC_{50} = 0.13 \mu M$, $EC_{50} = 1.4 \mu M$) analogs inhibit both NS5b (as the NTP) and the HCV replicon [65]. Mechanism of action work clearly demonstrates that these nucleosides act as chain-terminators. Modeling of 38 into the RNA primer portion of the NS5b crystal structure suggests how these 3'-hydroxyl containing nucleosides are able to act as chain-terminators. The 2'-methylribose nucleoside is incorporated as a substrate but the 2'-methyl group prevents subsequent incorporation via steric clashes with either the O4' or H8 of purines of the incoming nucleotide. Resistant replicons were generated against 38 and showed a S282T mutation in the NS5b sequence. When this mutation is modeled with 38 into the crystal structure, the addition of a methyl group to the wild-type serine (i.e., threonine) clearly prevents the incorporation of this nucleotide into the resistant mutant [65]. The cytidine analog (40) is claimed to be active *in vitro* as the active metabolite of its 3'-valyl ester NM-283 (41) [66,67] which is currently undergoing a phase Ib/phase IIa clinical trial. Initial data from a 50 patient study (doses from 25 to 800 mg) demonstrated an approximately 0.5 \log_{10} reduction in HCV RNA levels after 100 mg once-daily dosing for 15 days. Clinical pharmacokinetics at both day 1 and 15 showed a half-life slightly longer than 4 h with no apparent accumulation [66,68].



Recently, base modified analogs containing 2'-methylribose have been reported, including the tubercidin (42) and 6-thiophene-purine (43) [69,70]. Multiple modified-base nucleosides including a hydrazine analog series of 44 and thiomethylpurine analog 45 were reported to have HCV replicon EC_{50} s less than $10 \mu M$ [71-74]. Mouse PK data for 45 demonstrated half-lives of 1.2 h and 0.43 h for oral and IV dosing, respectively, with no toxicity observed up to 160 mg/kg.

2.5. Other viral targets

HCV IRES inhibitors are mostly nucleic acid based molecules and no small molecule with promising activity has been reported. The only inhibitor that has demonstrated clinical effectiveness is ISIS 14803 (anti-sense oligonucleotide). In a phase II trial involving 43 patients with chronic hepatitis C, ISIS 14803 reduced viral loads in 7 patients from ≥ 1 to $3.8 \log_{10}$ copies/ml during a 12-week treatment period [75]. Gene silencing by RNA interference is being explored as a new approach to inhibit HCV replication. Duplex small interfering RNAs directed at the HCV genome has been shown to be effective in dramatically inhibiting HCV replication and protein expression in human hepatoma cells [76]. *In vivo* application is likely to require gene delivery.

Despite certain efforts in targeting the HCV NS3 helicase, no compound that selectively and directly inhibits the NS3 ATP-hydrolyzing or nucleic acid-unwinding activities has been reported. Several nucleoside compounds including ring-expanded nucleosides 46–48 were synthesized and evaluated for their abilities to affect helicases of *Flaviviridae* viruses. Mixed results were obtained. The compounds can cause either enzyme inhibition or activation depending on the assay condition, source of the enzyme and nucleic acid template [77]. While these compounds may be interesting tools to elucidate putative allosteric sites on the helicase, their ability to inhibit HCV replication remains to be shown.

The HCV P7 protein forms an ion channel in lipid bilayers. It is not essential for viral replication, but it has been speculated that cation permeation across membranes may be important for the maturation and release of infectious virions. There is interest in evaluating P7 as an anti-viral target. Amantidine, 49, an anti-influenza agent that inhibits the ion channel of influenza virus, inhibits the HCV P7 ion channel [78]. Amantidine has been tested clinically against HCV and as a single agent has not shown anti-viral activity; in combination with interferon or interferon and ribavirin has yielded mixed results [79,80]. Long-alkyl-chain inosugar derivatives inhibit bovine viral diarrhea virus (BVDV). 50–52 were recently shown to inhibit the HCV P7 ion channel in black lipid membrane [81].

3. INHIBITORS FOR NON-VIRAL TARGETS

3.1. Caspase


Liver disease caused by conditions such as HCV infection is characterized by excessive apoptosis. Oxamyl dipeptide IDN-6556 is an anti-apoptotic irreversible caspase inhibitor with potencies against several caspases in the low to subnanomolar range [82]. In a phase IIa trial with chronic hepatitis C patients, oral dosing of IDN-6556 for 14 days significantly lowered liver enzyme levels in plasma at all doses [83]. While IDN-6556 is not an anti-viral agent, it may help infected patients to preserve the function and health of their livers.

3.2. Toll-like receptor 7

Certain C8-substituted and N7, C8-substituted guanosine analogs stimulate immune responses. Some of these compounds activate toll-like receptor 7 (TLR7), a member

HCV Anti-A

of a group
ANA245,
oral pro-c
2',5'-oligo
response.
reduced
maximal
with indu



3.3. Pro

A fortuito
in HCV I
dose-depe

3.4. CD

CD81 is a
to be a bi
enough fo
hepatocyt
Small mo
cells [87]
determine


R.C. Griffith et al.

HCV Anti-viral Agents

233

no small molecule it has demonstrated. In a phase II trial, reduced viral loads in 7 days period [75]. Gene therapy to inhibit HCV protein expression in gene delivery. Found that selectively binding activities has reduced nucleosides 46-48 uses of *Flaviviridae* reverse transcriptase and nucleic acid polymerase to elucidate putative mechanisms remains to be shown. It is essential for viral replication in membranes may be. There is interest in a agent that inhibits [8]. Amantadine has anti-viral activity; in reduced results [79,80]. Borna disease virus (BVDV). 50-nm lipid membrane [81].

of a group of receptors that take part in the host defense against viruses. Isatoribine (53, ANA245) is a TLR7 agonist that is currently under evaluation in the clinic and so is its oral pro-drug ANA971 [84]. While it has no direct anti-viral activity, it induces hepatic 2',5'-oligoadenylate synthetase (OAS) leading to activation of the innate immune response. In a phase Ib trial with a 7-day treatment by IV administration, the compound reduced viral load in patients with chronic HCV infection. Viral load reduction was maximal in the 800 mg arm at approximately 1-3 log₁₀ copies/ml and was associated with induction of the OAS system [85].



3.3. Proteasome inhibitors

A fortuitous discovery recently implicated the human 20S proteasome subunit α -PSMA7 in HCV IRES mediated translation. In fact, a proteasome inhibitor MG132 54 showed dose-dependent inhibition of HCV IRES activity in human hepatoma cells [86].

3.4. CD81

CD81 is a member of the tetraspanin family of integral membrane protein and is thought to be a binding partner for the HCV envelope E2 protein. Although binding alone is not enough for infection, prevention of binding does coincide with lack of infection in human hepatocytes. An extracellular region (D helix) on CD81 is important for binding to E2. Small molecule mimics 55, 56 of the D helix inhibit binding of E2 to CD81 on Molt-4 cells [87]. Cellular activities of these compounds are unknown and may be difficult to determine since they require 6% DMSO for solubilization.

erized by excessive caspase inhibitor [82]. In a phase 1 trial, 14 days significantly reduced viral load in their livers.

stimulate immune response (TLR7), a member

4. CONCLUSION

There has been a dramatic increase in patents and publications describing new HCV inhibitors last year as many research organizations have turned their attention to this important viral disease. Compounds which directly target viral specific proteins are beginning to enter clinical trials and the recent encouraging proof-of-principle results with BILN 2061 will hopefully mark the beginning of an era of new high efficacy treatments for HCV.

REFERENCES

- [1] B. Szabo, G. Lotz, C. Paska, A. Kiss and Z. Schaff, *Pathol. Oncol. Res.*, 2003, 9, 215.
- [2] W. R. Kim, J. B. Gross, Jr., J. J. Poterucha, G. R. Locke, III and B. R. Dickson, *Hepatology*, 2001, 33, 201.
- [3] G. Lake-Bakaar, *Curr. Drug Targets. Infect. Disord.*, 2003, 3, 247.
- [4] S. Rosenberg, *J. Mol. Biol.*, 2001, 313, 451.
- [5] L. Pacini, A. Vitelli, G. Filocamo, L. Bartholomew, M. Brunetti, A. Tramontano, C. Steinkuhler and G. Migliaccio, *J. Virol.*, 2000, 74, 10563.
- [6] A. Lahm, A. Yagnik, A. Tramontano and U. Koch, *Curr. Drug Targets*, 2002, 3, 281.
- [7] Y. S. Tsantrizos, G. Bolger, P. Bonneau, D. R. Cameron, N. Goudreau, G. Kukolj, S. R. LaPlante, M. Llinas-Brunet, H. Nar and D. Lamarre, *Angew. Chem. Int. Ed.*, 2003, 42, 1356.
- [8] D. Lamarre, P. C. Anderson, M. Bailey, P. Beaulieu, G. Bolger, P. Bonneau, M. Boes, D. R. Cameron, M. Cartier, M. G. Cordingley, A. M. Faucher, N. Goudreau, S. H. Kawai, G. Kukolj, L. Lagace, S. R. LaPlante, H. Narjes, M. A. Poupert, J. Rancourt, R. E. Sentjens, R. St. George, B. Simoneau, G. Steinmann, D. Thibeault, Y. S. Tsantrizos, S. M. Weldon, C. L. Yong and M. Llinas-Brunet, *Nature*, 2003, 426, 186.
- [9] Y. Benhamou, HEP DART 2003, Kauai, HI, December 14–18, Abstract 025, 2003.
- [10] Boehringer Ingelheim Pharmaceuticals, http://us.boehringer-ingelheim.com/about/pressreleases/102803_hepatitis.html (last accessed 02/10/04).
- [11] E. Foy, K. Li, C. Wang, R. Sumpter, Jr., M. Ikeda, S. M. Lemon and M. Gale, Jr., *Science*, 2003, 300, 1145.
- [12] M. Llinas-Brunet and V. J. Gorys, *PCT Patent Appl. WO 2003064455 A2*, 2003.
- [13] J. A. Campbell and A. C. Good, *PCT Patent Appl. WO 2003053349 A2*, 2003.
- [14] M. Llinas-Brunet and V. J. Gorys, *PCT Patent Appl. WO 2003064456 A1*, 2003.
- [15] X. A. Wang, L.-Q. Sun, S.-Y. Sit, N. Sin, P. M. Scola, P. Hewawasam, A. C. Good, Y. Chen and J. A. Campbell, *PCT Patent Appl. WO 2003099274 A1*, 2003.
- [16] A. Johansson, A. Poliakov, E. Akerblom, K. Wiklund, G. Lindeberg, S. Winiwarter, U. H. Danielson, B. Samuelsson and A. Hallberg, *Bioorg. Med. Chem.*, 2003, 11, 2551.
- [17] V. R. Hegde, H. Pu, M. Patel, P. R. Das, N. Butkiewicz, G. Arreaza, V. P. Gullo and T. M. Chan, *Bioorg. Med. Chem. Lett.*, 2003, 13, 2925.
- [18] R. B. Pemi, G. Chandorkar, P. R. Chaturvedi, L. F. Courtney, C. J. Decker, C. A. Gates, S. L. Harbeson, A. D. Kwong, C. Lin, Y. P. Luong, W. Markland, G. Rao, R. D. Tung and J. A. Thomson, 54th Annu. Meet. Am. Assoc. Study Liver Dis., October 24–28, Boston, MA, Abstract 972, 2003.
- [19] C. Lin, K. Lin, C. A. Gates, D. Ma, D. Brennan, J. Fulghum, H.-M. Hsiao, G. Rao, Y. Wei, J. Alford, R. B. Pemi and A. D. Kwong, 54th Annu. Meet. Am. Assoc. Study Liver Dis., October 24–28, Boston, MA, Abstract 1000, 2003.
- [20] J. Pitlik, K. Cotrell, L. Farmer, R. Pemi, J. Van Drie and M. Mureko, *PCT Patent Appl. WO 2003087092 A2*, 2003.
- [21] R. B. Pemi, J. Pitlik, J. H. Van Drie and L. J. Farmer, *PCT Patent Appl. WO 2003035060 A1*, 2003.
- [22] F. Victor, J. Lamar, N. Snyder, Y. Yip, D. Guo, N. Yumibe, R. B. Johnson, Q. M. Wang, J. I. Glass and S. H. Chen, *Bioorg. Med. Chem. Lett.*, 2004, 14, 257.
- [23] W. Han, Z. Hu, X. Ji, 13, 1111.
- [24] R. B. Pemi, S. D. Bri, S. L. Harbeson, J. L. B. G. Rao, W. C. Sc Med. Chem. Lett., 21
- [25] D. M. Andrews, S. J. G. Mills, J. E. Robir
- [26] M. J. Slater, E. M. A Jones, G. Mills, N. R. J
- [27] D. M. Andrews, H. M P. S. Jones, G. Mills Eur. J. Med. Chem.,
- [28] J. Z. Wu and Z. Hor
- [29] D. O'Farrell, R. Tro
- [30] J. Schmidt-Mende, F J. Biol. Chem., 2001
- [31] G. McKercher, P. L. and G. Kukolj, *Nucl*
- [32] S. Bressanelli, L. To
- [33] C. A. Lesburg, R. R
- [34] H. Hashimoto, K. M
- [35] L. Tomei, S. Altam N. Gennari, M. Bist Francesco and G. M
- [36] D. Latour, C. Hanc J. J. Pouliot, HEP D
- [37] P. L. Beaulieu, G. F. *PCT Patent Appl. W*
- [38] P. L. Beaulieu, G. F J. J. Pouliot, HEP D
- [39] P. L. Beaulieu, G. F. *Appl. WO 03007945*
- [40] E. S. Priestly, C. P. I A2, 2003.
- [41] G. W. Shipp Jr., K. *Patent Appl. WO 03*
- [42] R. A. Love, H. B. F L. Wang, J. A. Thon
- [43] M. Wang, K. K. Ng N. Nguyen-Ba, M. F 278, 9489.
- [44] L. Chan, T. J. Red N. Nguyen-Ba, L. M. Q. Zhang, M. D M. Wang, M. M. C Abstract 124, 2003.
- [45] A. J. Borchardt and
- [46] J. Gonzalez, A. J. Bc H. Tatlock, M. A. A
- [47] G. Migliaccio, 16th
- [48] T. T. Nguyen, A. T Sarisky, *Antimicrob.*
- [49] C. Gardelli, C. Giul S. Ponzi, I. Stansfield

BEST AVAILABLE COPY

- [23] W. Han, Z. Hu, X. Jiang, Z. R. Wasserman and C. P. Decicco, *Bioorg. Med. Chem. Lett.*, 2003, 13, 1111.
- [24] R. B. Perni, S. D. Britt, J. C. Court, L. F. Courtney, D. D. Deininger, L. J. Farmer, C. A. Gats, S. L. Harbeson, J. L. Kim, J. A. Landro, R. B. Levin, Y. P. Luong, B. T. O'Malley, J. Pitluk, B. G. Rao, W. C. Schairer, J. A. Thomson, R. D. Tung, J. H. Van Drie and Y. Wei, *Bioorg. Med. Chem. Lett.*, 2003, 13, 4059.
- [25] D. M. Andrews, S. J. Carey, H. Chaignot, B. A. Coomber, N. M. Gray, S. L. Hind, P. S. Jones, G. Mills, J. E. Robinson and M. J. Slater, *Org. Lett.*, 2002, 4, 4475.
- [26] M. J. Slater, B. M. Amphlett, D. M. Andrews, P. Bamborough, S. J. Carey, M. R. Johnson, P. S. Jones, G. Mills, N. R. Parry, D. O. Somers, A. J. Stewart and T. Skarzynski, *Org. Lett.*, 2003, 5, 4627.
- [27] D. M. Andrews, H. M. Chaignot, B. A. Coomber, M. D. Dowle, S. Lucy Hind, M. R. Johnson, P. S. Jones, G. Mills, A. Patikis, T. J. Pateman, J. E. Robinson, M. J. Slater and N. Trivedi, *Eur. J. Med. Chem.*, 2003, 38, 339.
- [28] J. Z. Wu and Z. Hong, *Curr. Drug Targets. Infect. Disord.*, 2003, 3, 207.
- [29] D. O'Farrell, R. Trowbridge, D. Rowlands and J. Jager, *J. Mol. Biol.*, 2003, 326, 1025.
- [30] J. Schmidt-Mende, E. Bieck, T. Hagle, F. Penin, C. M. Rice, H. E. Blum and D. Moradpour, *J. Biol. Chem.*, 2001, 276, 44052.
- [31] G. McKercher, P. L. Beaulieu, D. Lamarre, S. LeFebvre, C. Pellerin, L. Thauvette and G. Kukolj, *Nucl. Acids Res.*, 2004, 32, 422.
- [32] S. Bressanelli, L. Tomei, F. A. Rey and R. De Francesco, *J. Virol.*, 2002, 76, 3492.
- [33] C. A. Lesburg, R. Radfar and P. C. Weber, *Curr. Opin. Invest. Drugs*, 2000, 1, 289.
- [34] H. Hashimoto, K. Mizutani and A. Yoshida, *Eur. Patent Appl.* EP1162196 A1, 2001.
- [35] L. Tomei, S. Altamura, L. Bartholomew, A. Biroccio, A. Ceccacci, L. Pacini, F. Narjes, N. Gennari, M. Bisocchi, I. Incitti, L. Orsatti, S. Harper, I. Stansfield, M. Rowley, R. De Francesco and G. Migliaccio, *J. Virol.*, 2003, 77, 13225.
- [36] D. Latour, C. Hancock, E. Michelotti, L. Lou, D.-F. Shi, C. D. Roberts, R. Griffith and J. J. Pouliot, HEP DART 2003, Kanai, HI, December 14-18, Abstract 032, 2003.
- [37] P. L. Beaulieu, G. Fazal, G. Kukolj, E. Jolicoeur, J. Gillard, M. A. Poupart and J. Rancourt, *PCT Patent Appl.* WO 03010140 A2, 2003.
- [38] P. L. Beaulieu, G. Fazal, S. Goulet, G. Kukolj, M. Poirier, Y. S. Tsantrizos, E. Jolicoeur, J. Gillard, M. A. Poupart and J. Rancourt, *PCT Patent Appl.* WO 03010141 A2, 2003.
- [39] P. L. Beaulieu, G. Fazal, S. Goulet, G. Kukolj, M. Poirier and Y. S. Tsantrizos, *PCT Patent Appl.* WO 03007945 A1, 2003.
- [40] E. S. Priestly, C. P. Decicco, T. W. Hudyma and X. Zheng, *PCT Patent Appl.* WO 03026587 A2, 2003.
- [41] G. W. Shipp Jr., K. E. Rosner, J. Popovici-Muller, Y. Deng, T. Wang and P. J. Curran, *PCT Patent Appl.* WO 03101993 A1, 2003.
- [42] R. A. Love, H. E. Parge, X. Yu, M. J. Hickey, W. Diehl, J. Gao, H. Wriggers, A. Ekker, L. Wang, J. A. Thomson, P. S. Dragovich and S. A. Fuhrman, *J. Virol.*, 2003, 77, 7575.
- [43] M. Wang, K. K. Ng, M. M. Cherney, L. Chan, C. G. Yannopoulos, J. Bedard, N. Morin, N. Nguyen-Ba, M. H. Alaoui-Ismaili, R. C. Bethell and M. N. James, *J. Biol. Chem.*, 2003, 278, 9489.
- [44] L. Chan, T. J. Reddy, M. Proulx, S. K. Das, O. Pereira, C. Yannopoulos, C. Poisson, N. Nguyen-Ba, L. Halab, M. Courchesne, C. Roy, R. Bethell, A. Siddiqui, W. Wang, M. Q. Zhang, M. David, M. Hamel, L. L'Heureux, J. Bedard, O. Nicolas, M. N. James, M. Wang, M. M. Cherney and K. K. Ng, 16th ICAR, Savannah, GA, April 27-May 1, Abstract 124, 2003.
- [45] A. J. Borchardt and M. P. Goble, *PCT Patent Appl.* WO 03082848 A1, 2003.
- [46] J. Gonzalez, A. J. Borchardt, P. S. Dragovich, T. M. Jewell, M. A. Linton, R. Zhou, H. Li, J. H. Tatlock, M. A. Abreo and T. J. Prins, *PCT Patent Appl.* WO 03095441 A1, 2003.
- [47] G. Migliaccio, 16th ICAR, Savannah, GA, April 27-May 1, LB-12, 2004.
- [48] T. T. Nguyen, A. T. Gats, L. L. Gutshall, V. K. Johnston, B. Gu, K. J. Duffy and R. T. Sarisky, *Antimicrob. Agents Chemother.*, 2003, 47, 3525.
- [49] C. Gardelli, C. Giuliano, S. Harper, U. Koch, F. Narjes, J. M. Ontoria Ontoria, M. Poma, S. Ponzi, I. Stansfield and V. Summa, *PCT Patent Appl.* WO 0206246 A1, 2002.

BEST AVAILABLE COPY

- [50] Viropharma, http://www.irconnect.com/vphm/pages/news_releases.html?d=42687 (last accessed 03/26/04).
- [51] A. Gopalsamy, J. W. Ellingboe and T. S. Mansour, *PCT Patent Appl.* WO 03099824 A1, 2003.
- [52] A. Gopalsamy, M. S. Collette, J. W. Ellingboe, M. G. Laporte, T. S. Mansour, C. J. Burns, K. Park and S. M. Condon, *PCT Patent Appl.* WO 03099275 A1, 2003.
- [53] D. Goff, H. Lu, R. Singh, T. Sun and S. D. Issankani, *PCT Patent Appl.* WO 03040112, 2003.
- [54] Rigel Pharmaceuticals, http://www.rigel.com/genomics/pr_1074900710 (last accessed 03/26/04).
- [55] G. Bravi, H. S. Goodland, D. Haigh, C. D. Hartley, V. L. H. Lovegrove, P. Shah and M. J. Slater, *PCT Patent Appl.* WO 03037894 A1, 2003.
- [56] G. Bravi, R. Guidetti, D. Haigh, C. D. Hartley, P. D. Howes, D. L. Jackson, V. L. H. Lovegrove, P. Shah, M. J. Slater and K. J. Wareing, *PCT Patent Appl.* WO 03037895 A1, 2003.
- [57] G. Bravi, P. D. Howes, V. L. H. Lovegrove, P. Shah and M. J. Slater, *PCT Patent Appl.* WO 03037893 A1, 2003.
- [58] Y. H. Chung, C. W. Lee, W. Y. Yang, H. D. Lee, W. J. Park, Y. Y. Chang and J. S. Lee, *PCT Patent Appl.* WO 02079187 A1, 2002.
- [59] Y. H. Chung, C. W. Lee, W. Y. Yang, H. D. Lee, W. J. Park, Y. Y. Chang and J. S. Lee, *PCT Patent Appl.* WO 03002518 A1, 2003.
- [60] H. Gao, M. L. Greene, R. J. Gross, R. A. Nugent and J. Pfeifferkorn, *PCT Patent Appl.* WO 2004002977 A1, 2004.
- [61] B. C. Finzel, L. A. Funk, R. C. Kelly, M. T. Reding and N. A. Wicnienski, *PCT Patent Appl.* WO 2004002944 A1, 2004.
- [62] R. R. Devos, *PCT Patent Appl.* 6,660,721 B2, 2003.
- [63] R. R. Devos, *PCT Patent Appl.* WO 2002100415 A2, 2002.
- [64] R. Storer, *PCT Patent Appl.* WO 200132153 A2, 2001.
- [65] G. Migliaccio, J. E. Tomassini, S. S. Carroll, L. Tomci, S. Altamura, B. Bhat, L. Bartholomew, M. R. Bosserman, A. Ceccacci, L. F. Colwell, R. Cortese, R. De Francesco, A. B. Eldrup, K. L. Getty, X. S. Hou, R. L. LaFemina, S. W. Ludmerer, M. MacCoss, D. R. McMaster, M. W. Stahlhut, D. B. Olsen, D. J. Hazuda and O. A. Flores, *J. Biol. Chem.*, 2003, 278, 49164.
- [66] J.-P. Sommadossi, *PCT Patent Appl.* WO 2004002422 A2, 2004.
- [67] J.-P. Sommadossi, *PCT Patent Appl.* WO 2004003000 A2, 2004.
- [68] Idenix Pharmaceuticals, <http://www.sec.gov/Archives/edgar/data/1093649/000095013503006012/0000950135-03-006012.txt>.
- [69] S. S. Carroll, J. E. Tomassini, M. Bosserman, K. Getty, M. W. Stahlhut, A. B. Eldrup, B. Bhat, D. Hall, A. L. Simcoe, R. LaFemina, C. A. Rutkowski, B. Wolanski, Z. Yang, G. Migliaccio, R. De Francesco, L. C. Kuo, M. MacCoss and D. B. Olsen, *J. Biol. Chem.*, 2003, 278, 11979.
- [70] S. S. Carroll, R. L. LaFemina, D. L. Hall, A. L. Himmelberger, L. C. Kuo, M. MacCoss, D. B. Olsen, C. A. Rutkowski, J. E. Tomassini, H. An, B. Bhat, N. Bhat, P. D. Cook, A. B. Eldrup, C. J. Guinosso, M. Prihac and T. P. Prakash, *PCT Patent Appl.* 02057425 WO, 2002.
- [71] H. An and Y. Ding, *PCT Patent Appl.* WO 2003061576 A2, 2003.
- [72] H. An, Y. Ding, S. Shaw and Z. Hong, *PCT Patent Appl.* WO 03062256 A1, 2003.
- [73] Z. Hong, H. An, Y. Ding, J. L. Girardet and W. Zhong, *PCT Patent Appl.* WO 03062255 A2, 2003.
- [74] H. An, Z. Hong, K. Smith, Y. Ding and J. L. Girardet, *PCT Patent Appl.* WO 03061385 A1, 2003.
- [75] S. C. Gordon, B. R. Bacon, I. M. Jacobson, L. Shiffman, N. H. Afdhal, R. Z. Yu, J. G. McHutchison and J. K. Kwoh, 54th Annu. Meet. Am. Assoc. Study Liver Dis., October 24-28, Boston, MA, Abstract 312, 2003.
- [76] T. Yokota, N. Sakamoto, N. Enomoto, Y. Tanabe, M. Miyagashi, S. Maekawa, M. Kurosaki, L. Yi, K. Taira, M. Watanabe and H. Mizusawa, *EMBO Rep.*, 2003, 4, 602.

- [77] N. Zhan
Fatom,
- [78] S. D. Gr
and S. I
- [79] G. Ideo
A. Fran
M. Sarri
Abstract
- [80] E. J. La
A. Gasin
Meet. A
- [81] D. Pavl
N. Zito
- [82] D. S. K
- [83] D. Shap
M. Huy
087, 20
- [84] J. Lee, T.
2003, 1
- [85] B. Kerr
J. Theis
DART,
- [86] M. Kru
Biol., 2
- [87] S. E. V.
Todd, 1

R.C. Griffith et al.

al?d = 42687 (last

WO 03099824 A1,

ansour, C. J. Burns,

VO 03040112, 2003.

710 (last accessed

Shah and M. J. Slater,

Jackson, V. L. H.

WO 03037895 A1,

CT Patent Appl. WO

ng and J. S. Lee, PCT

ng and J. S. Lee, PCT

CT Patent Appl. WO

ski, PCT Patent Appl.

Itamura, B. Bhat, L.
ese, R. De Francesco,
er, M. MacCoss, D. R.
i, J. Biol. Chem., 2003,

J1093649/000095013

tahlhut, A. B. Eldrup,
3. Wolanski, Z. Yang,
Olsen, J. Biol. Chem.,

C. Kuo, M. MacCoss,
Bhat, P. D. Cook, A. B.
4. 02057425 WO, 2002.

2256 A1, 2003.

pp1. WO 03062255 A2,

pp1. WO 03061385 A1,

H. Afdhal, R. Z. Yu,
tudy Liver Dis., October

Maekawa, M. Kurosaki,
, 4. 602.

HCV Anti-viral Agents

237

- [77] N. Zhang, H. M. Chen, V. Koch, H. Schmitz, C. L. Liao, M. Bretner, V. S. Bharti, A. I. Patton, R. B. Naso, R. S. Hosmane and P. Borowski, *J. Med. Chem.*, 2003, 46, 4149.
- [78] S. D. Griffin, D. J. Rowlands, D. S. Clarke, J. Jaeger, L. P. Beales, M. P. Harris, O. Worsfold and S. D. Evans, *FEBS Lett.*, 2003, 535, 34.
- [79] G. Ideo, G. Antonucci, A. F. Atilli, S. Babudieri, A. Chirianai, A. Craxi, G. DiPerri, A. Francavilla, A. Mangia, A. Piccotto, M. Rizzetto, G. Ruggiero, F. Suter, S. Tripi and M. Sarracino, 54th Annu. Meet. Am. Assoc. Study Liver Dis., October 24-28, Boston, MA, Abstract 1186, 2003.
- [80] E. J. Lawitz, N. S. Cantu, M. Davis, N. Afdahl, M. Curry, M. Maillard, F. Adams, N. Tsai, A. Gasic, T. Kothari, J. Cox, D. P. Ganeshappa, B. Silverman and S. C. Kadakia, 54th Annu. Meet. Am. Assoc. Study Liver Dis., October 24-28, Boston, MA, Abstract 997, 2003.
- [81] D. Pavlovic, D. C. A. Neville, O. Argaud, B. Blumberg, R. A. Dwek, W. B. Fischer and N. Zitzmann, *PNAS*, 2003, 100, 6104.
- [82] D. S. Karanewsky and R. J. Teranosky, *PCT Patent Appl.* US 6,199,750, 2001.
- [83] D. Shapiro, E. Schiff, P. Pockros, M. Shiffman, J. McHutchison, A. Muir, R. Gish, N. Afdahl, M. Huyghe, D. Hecht and T. Oltersdorf, HEP DART, December 14-18, Kauli, HI, Abstract 087, 2003.
- [84] J. Lee, T. H. Chuang, V. Redecke, P. M. Pitha, D. A. Carson, E. Raz and H. B. Cottam, *PNAS*, 2003, 100, 6646.
- [85] B. Kerr, L. Bauman, W. Liaw, S. Webber, L. Kirkovaky, K. Steffy, S. Fletcher, V. Banh, J. Theiss, Y. Horsmans, J. P. Desager, T. Berg, T. Mueller, S. Worland and D. Averett, HEP DART, December 14-18, Kauli, HI, Abstract 063, 2003.
- [86] M. Kruger, C. Berger, P. J. Welch, J. R. Barber, M. P. Manns and F. Wong-Staal, *Mol. Cell. Biol.*, 2001, 24, 8657.
- [87] S. E. VanCompernelle, A. V. Wiznycia, J. R. Rush, M. Dhanasekaran, P. W. Baures and S. C. Todd, *Virology*, 2003, 314, 371.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.